

3.2.2 死因

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=2861	BU(経口)を含むレジメン n=317	
拒絶・生着不全	70 (2.5%)	11 (3.5%)	P = 0.03
間質性肺炎以外の感染症	557 (19.5%)	49 (15.5%)	
間質性肺炎	153 (5.3%)	20 (6.3%)	
ARDS	43 (1.5%)	4 (1.3%)	
急性GVHD	137 (4.8%)	16 (5.1%)	
慢性GVHD	61 (2.1%)	20 (6.3%)	
VOD	79 (2.8%)	9 (2.8%)	
TMA	79 (2.8%)	10 (3.2%)	
原疾患の再発・増悪	1106 (38.7%)	112 (35.3%)	
二次性悪性腫瘍	16 (0.6%)	1 (0.3%)	
出血	151 (5.3%)	11 (3.5%)	
外因死(自殺・事故含む)	5 (0.2%)	2 (0.6%)	
臓器不全(上記以外の原因による)	282 (9.9%)	41 (13.0%)	
その他	122 (4.3%)	11 (3.5%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=655	CY+TBI ±other drugs n=2433	BU(経静脈) +FL±other drugs n=2091	FL+TBI ±other drugs n=2168	
拒絶・生着不全	16 (2.4%)	55 (2.3%)	49 (2.3%)	95 (4.4%)	P < 0.001
間質性肺炎以外の感染症	98 (15.0%)	419 (17.2%)	440 (21.1%)	462 (21.3%)	
間質性肺炎	31 (4.7%)	161 (6.6%)	117 (5.6%)	111 (5.1%)	
ARDS	10 (1.5%)	45 (1.9%)	32 (1.5%)	40 (1.9%)	
急性GVHD	31 (4.7%)	87 (3.6%)	98 (4.7%)	86 (4.0%)	
慢性GVHD	18 (2.8%)	73 (3.0%)	43 (2.1%)	35 (1.6%)	
VOD	22 (3.4%)	85 (3.5%)	54 (2.6%)	27 (1.3%)	
TMA	16 (2.4%)	59 (2.4%)	61 (2.9%)	59 (2.7%)	
原疾患の再発・増悪	276 (42.1%)	1009 (41.5%)	776 (37.1%)	749 (34.6%)	
二次性悪性腫瘍	0 (0.0%)	10 (0.4%)	16 (0.8%)	16 (0.7%)	
出血	29 (4.4%)	95 (3.9%)	118 (5.6%)	107 (4.9%)	
外因死(自殺・事故含む)	4 (0.6%)	6 (0.3%)	1 (0.1%)	5 (0.2%)	
臓器不全(上記以外の原因による)	57 (8.7%)	227 (9.3%)	215 (10.3%)	255 (11.8%)	
その他	47 (7.2%)	102 (4.2%)	71 (3.4%)	121 (5.6%)	