

3.4.2 性腺機能障害

男性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1454	BU(経口)を含むレジメン n=152	
性腺機能障害無	198 (13.6%)	19 (12.5%)	$P = 0.98$
性腺機能障害有	29 (2.0%)	3 (2.0%)	
未検査	793 (54.5%)	85 (55.9%)	
不明	434 (29.9%)	45 (29.6%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=468	CY+TBI±other drugs n=1911	BU(経静脈)+FL±other drugs n=913	FL+TBI±other drugs n=1063	
性腺機能障害無	59 (12.6%)	229 (12.0%)	127 (13.9%)	167 (15.7%)	$P = 0.004$
性腺機能障害有	8 (1.7%)	38 (2.0%)	19 (2.1%)	9 (0.9%)	
未検査	267 (57.1%)	1138 (59.6%)	482 (52.8%)	585 (55.0%)	
不明	134 (28.6%)	506 (26.5%)	285 (31.2%)	302 (28.4%)	

女性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1087	BU(経口)を含むレジメン n=111	
性腺機能障害無	112 (10.3%)	18 (16.2%)	$P = 0.25$
性腺機能障害有	93 (8.6%)	11 (9.9%)	
未検査	557 (51.2%)	52 (46.9%)	
不明	325 (29.9%)	30 (27.0%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=342	CY+TBI±other drugs n=1526	BU(経静脈)+FL±other drugs n=686	FL+TBI±other drugs n=757	
性腺機能障害無	31 (9.1%)	157 (10.3%)	74 (10.8%)	111 (14.7%)	$P < 0.001$
性腺機能障害有	34 (9.9%)	330 (21.6%)	51 (7.4%)	71 (9.4%)	
未検査	179 (52.3%)	693 (45.4%)	352 (51.3%)	381 (50.3%)	
不明	98 (28.7%)	346 (22.7%)	209 (30.5%)	194 (25.6%)	