

3.2.2 死因

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=3,581	BU(経口)を含むレジメン n=340	
拒絶・生着不全	76 (2.1%)	13 (3.8%)	P = 0.01
間質性肺炎以外の感染症	675 (18.9%)	50 (14.7%)	
間質性肺炎	192 (5.4%)	21 (6.2%)	
ARDS	53 (1.5%)	4 (1.2%)	
急性GVHD	172 (4.8%)	16 (4.7%)	
慢性GVHD	83 (2.3%)	22 (6.5%)	
VOD	102 (2.9%)	9 (2.7%)	
TMA	103 (2.9%)	12 (3.5%)	
原疾患の再発・増悪	1,395 (39.0%)	121 (35.6%)	
二次性悪性腫瘍	23 (0.6%)	1 (0.3%)	
出血	171 (4.8%)	11 (3.2%)	
外因死(自殺・事故含む)	7 (0.2%)	2 (0.6%)	
臓器不全(上記以外の原因による)	363 (10.1%)	45 (13.2%)	
その他	166 (4.6%)	13 (3.8%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=800	CY+TBI ±other drugs n=2,849	BU(経静脈) +FL±other drugs n=2,642	FL+TBI ±other drugs n=2,690	
拒絶・生着不全	17 (2.1%)	60 (2.1%)	54 (2.0%)	107 (4.0%)	P < .001
間質性肺炎以外の感染症	113 (14.1%)	489 (17.2%)	540 (20.4%)	559 (20.8%)	
間質性肺炎	37 (4.6%)	183 (6.4%)	150 (5.7%)	138 (5.1%)	
ARDS	12 (1.5%)	53 (1.9%)	40 (1.5%)	48 (1.8%)	
急性GVHD	37 (4.6%)	101 (3.6%)	127 (4.8%)	113 (4.2%)	
慢性GVHD	22 (2.8%)	84 (3.0%)	60 (2.3%)	48 (1.8%)	
VOD	29 (3.6%)	99 (3.5%)	70 (2.7%)	32 (1.2%)	
TMA	26 (3.3%)	76 (2.7%)	75 (2.8%)	69 (2.6%)	
原疾患の再発・増悪	341 (42.6%)	1,163 (40.8%)	985 (37.3%)	941 (35.0%)	
二次性悪性腫瘍	2 (0.3%)	19 (0.7%)	20 (0.8%)	27 (1.0%)	
出血	34 (4.3%)	114 (4.0%)	133 (5.0%)	119 (4.4%)	
外因死(自殺・事故含む)	4 (0.5%)	8 (0.3%)	3 (0.1%)	6 (0.2%)	
臓器不全(上記以外の原因による)	71 (8.9%)	271 (9.5%)	278 (10.5%)	328 (12.2%)	
その他	55 (6.9%)	129 (4.5%)	107 (4.1%)	155 (5.8%)	