

### 3.4.2 性腺機能障害

#### 男性

|         | カテゴリ区分①                   |                        | 検定     |
|---------|---------------------------|------------------------|--------|
|         | BU(経静脈)を含むレジメン<br>n=1,792 | BU(経口)を含むレジメン<br>n=166 |        |
| 性腺機能障害無 | 250 (14.0%)               | 18 (10.8%)             | p=0.66 |
| 性腺機能障害有 | 31 (1.7%)                 | 4 (2.4%)               |        |
| 未検査     | 996 (55.6%)               | 94 (56.6%)             |        |
| 不明      | 515 (28.7%)               | 50 (30.1%)             |        |

|         | カテゴリ区分②                         |                               |                                   |                               | 検定     |
|---------|---------------------------------|-------------------------------|-----------------------------------|-------------------------------|--------|
|         | BU(経静脈)+CY±other drugs<br>n=550 | CY+TBI±other drugs<br>n=2,285 | BU(経静脈)+FL±other drugs<br>n=1,153 | FL+TBI±other drugs<br>n=1,349 |        |
| 性腺機能障害無 | 72 (13.1%)                      | 285 (12.5%)                   | 163 (14.1%)                       | 209 (15.5%)                   | p=0.01 |
| 性腺機能障害有 | 7 (1.3%)                        | 53 (2.3%)                     | 22 (1.9%)                         | 17 (1.3%)                     |        |
| 未検査     | 320 (58.2%)                     | 1,354 (59.3%)                 | 625 (54.2%)                       | 744 (55.2%)                   |        |
| 不明      | 151 (27.5%)                     | 593 (26.0%)                   | 343 (29.7%)                       | 379 (28.1%)                   |        |

#### 女性

|         | カテゴリ区分①                   |                        | 検定     |
|---------|---------------------------|------------------------|--------|
|         | BU(経静脈)を含むレジメン<br>n=1,343 | BU(経口)を含むレジメン<br>n=117 |        |
| 性腺機能障害無 | 147 (11.0%)               | 17 (14.5%)             | p=0.48 |
| 性腺機能障害有 | 115 (8.6%)                | 12 (10.3%)             |        |
| 未検査     | 705 (52.5%)               | 54 (46.2%)             |        |
| 不明      | 376 (28.0%)               | 34 (29.1%)             |        |

|         | カテゴリ区分②                         |                               |                                 |                             | 検定      |
|---------|---------------------------------|-------------------------------|---------------------------------|-----------------------------|---------|
|         | BU(経静脈)+CY±other drugs<br>n=413 | CY+TBI±other drugs<br>n=1,769 | BU(経静脈)+FL±other drugs<br>n=862 | FL+TBI±other drugs<br>n=979 |         |
| 性腺機能障害無 | 50 (12.1%)                      | 180 (10.2%)                   | 90 (10.4%)                      | 141 (14.4%)                 | p<0.001 |
| 性腺機能障害有 | 38 (9.2%)                       | 385 (21.8%)                   | 63 (7.3%)                       | 90 (9.2%)                   |         |
| 未検査     | 217 (52.5%)                     | 808 (45.7%)                   | 459 (53.3%)                     | 507 (51.8%)                 |         |
| 不明      | 108 (26.2%)                     | 396 (22.4%)                   | 250 (29.0%)                     | 241 (24.6%)                 |         |