

### 3.4.2 性腺機能障害

#### 男性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1,792	BU(経口)を含むレジメン n=166	
性腺機能障害無	250 (14.0%)	18 (10.8%)	p=0.66
性腺機能障害有	31 (1.7%)	4 (2.4%)	
未検査	996 (55.6%)	94 (56.6%)	
不明	515 (28.7%)	50 (30.1%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=550	CY+TBI±other drugs n=2,285	BU(経静脈)+FL±other drugs n=1,153	FL+TBI±other drugs n=1,349	
性腺機能障害無	72 (13.1%)	285 (12.5%)	163 (14.1%)	209 (15.5%)	p=0.01
性腺機能障害有	7 (1.3%)	53 (2.3%)	22 (1.9%)	17 (1.3%)	
未検査	320 (58.2%)	1,354 (59.3%)	625 (54.2%)	744 (55.2%)	
不明	151 (27.5%)	593 (26.0%)	343 (29.7%)	379 (28.1%)	

#### 女性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1,343	BU(経口)を含むレジメン n=117	
性腺機能障害無	147 (11.0%)	17 (14.5%)	p=0.48
性腺機能障害有	115 (8.6%)	12 (10.3%)	
未検査	705 (52.5%)	54 (46.2%)	
不明	376 (28.0%)	34 (29.1%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=413	CY+TBI±other drugs n=1,769	BU(経静脈)+FL±other drugs n=862	FL+TBI±other drugs n=979	
性腺機能障害無	50 (12.1%)	180 (10.2%)	90 (10.4%)	141 (14.4%)	p<0.001
性腺機能障害有	38 (9.2%)	385 (21.8%)	63 (7.3%)	90 (9.2%)	
未検査	217 (52.5%)	808 (45.7%)	459 (53.3%)	507 (51.8%)	
不明	108 (26.2%)	396 (22.4%)	250 (29.0%)	241 (24.6%)	