

3.2.2 死因

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=4,354	BU(経口)を含むレジメン n=355	
拒絶・生着不全	89 (2.0%)	13 (3.7%)	p=0.001
間質性肺炎以外の感染症	823 (18.9%)	54 (15.2%)	
間質性肺炎	227 (5.2%)	22 (6.2%)	
ARDS	61 (1.4%)	4 (1.1%)	
急性GVHD	214 (4.9%)	18 (5.1%)	
慢性GVHD	102 (2.3%)	22 (6.2%)	
VOD	125 (2.9%)	10 (2.8%)	
TMA	124 (2.9%)	12 (3.4%)	
原疾患の再発・増悪	1,698 (39.0%)	126 (35.5%)	
二次性悪性腫瘍	33 (0.8%)	3 (0.9%)	
出血	192 (4.4%)	11 (3.1%)	
外因死(自殺・事故含む)	9 (0.2%)	2 (0.6%)	
臓器不全(上記以外の原因による)	448 (10.3%)	46 (13.0%)	
その他	209 (4.8%)	12 (3.4%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=915	CY+TBI ±other drugs n=3,173	BU(経静脈) +FL±other drugs n=3,274	FL+TBI ±other drugs n=3,149	
拒絶・生着不全	19 (2.1%)	68 (2.1%)	65 (2.0%)	118 (3.8%)	p<0.001
間質性肺炎以外の感染症	128 (14.0%)	542 (17.1%)	666 (20.3%)	656 (20.8%)	
間質性肺炎	41 (4.5%)	197 (6.2%)	180 (5.5%)	165 (5.2%)	
ARDS	13 (1.4%)	57 (1.8%)	47 (1.4%)	57 (1.8%)	
急性GVHD	43 (4.7%)	112 (3.5%)	163 (5.0%)	130 (4.1%)	
慢性GVHD	24 (2.6%)	91 (2.9%)	76 (2.3%)	56 (1.8%)	
VOD	33 (3.6%)	107 (3.4%)	89 (2.7%)	39 (1.2%)	
TMA	31 (3.4%)	85 (2.7%)	91 (2.8%)	92 (2.9%)	
原疾患の再発・増悪	378 (41.3%)	1,316 (41.5%)	1,238 (37.8%)	1,104 (35.1%)	
二次性悪性腫瘍	2 (0.2%)	22 (0.7%)	30 (0.9%)	34 (1.1%)	
出血	36 (3.9%)	122 (3.8%)	152 (4.6%)	133 (4.2%)	
外因死(自殺・事故含む)	6 (0.7%)	9 (0.3%)	3 (0.1%)	6 (0.2%)	
臓器不全(上記以外の原因による)	89 (9.7%)	300 (9.5%)	341 (10.4%)	376 (11.9%)	
その他	72 (7.9%)	145 (4.6%)	133 (4.1%)	183 (5.8%)	