

3.4.2 性腺機能障害

男性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=2,158	BU(経口)を含むレジメン n=173	
性腺機能障害無	301 (14.0%)	21 (12.1%)	p=0.87
性腺機能障害有	38 (1.8%)	4 (2.3%)	
未検査	1,214 (56.3%)	100 (57.8%)	
不明	605 (28.0%)	48 (27.8%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=641	CY+TBI±other drugs n=2,607	BU(経静脈)+FL±other drugs n=1,413	FL+TBI±other drugs n=1,564	
性腺機能障害無	86 (13.4%)	328 (12.6%)	197 (13.9%)	236 (15.1%)	p=0.003
性腺機能障害有	7 (1.1%)	64 (2.5%)	28 (2.0%)	22 (1.4%)	
未検査	381 (59.4%)	1,556 (59.7%)	772 (54.6%)	874 (55.9%)	
不明	167 (26.1%)	659 (25.3%)	416 (29.4%)	432 (27.6%)	

女性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1,595	BU(経口)を含むレジメン n=121	
性腺機能障害無	164 (10.3%)	16 (13.2%)	p=0.67
性腺機能障害有	149 (9.3%)	13 (10.7%)	
未検査	834 (52.3%)	58 (47.9%)	
不明	448 (28.1%)	34 (28.1%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=483	CY+TBI±other drugs n=1,977	BU(経静脈)+FL±other drugs n=1,038	FL+TBI±other drugs n=1,160	
性腺機能障害無	53 (11.0%)	200 (10.1%)	102 (9.8%)	173 (14.9%)	p<0.001
性腺機能障害有	56 (11.6%)	433 (21.9%)	78 (7.5%)	106 (9.1%)	
未検査	252 (52.2%)	901 (45.6%)	550 (53.0%)	607 (52.3%)	
不明	122 (25.3%)	443 (22.4%)	308 (29.7%)	274 (23.6%)	